



Action Steps for Parents of Children at High Risk for Flu Complications

August 7, 2009 9:00 AM ET

Children with chronic health problems such as asthma, diabetes, heart disease, metabolic conditions, neurologic and neuromuscular disorders, or who are pregnant are at higher risk of having complications from flu. In addition, all children younger than 5 years old are at higher risk of flu complications compared to older children. If you are not sure if any of your children are at higher risk for flu complications, please check with a doctor.

Keep children at high risk for flu complications from getting sick with the flu

Make sure your child's hands are washed for 20 seconds with soap and water or an alcohol-based hand rub often and especially after coughing or sneezing.

Have your child **cough and sneeze into a tissue** or into his or her elbow or shoulder if a tissue is not available.

Keep your child away from people who are sick.

Clean surfaces and objects that your child frequently touches with cleaning agents that are usually used.

When there is flu in your community, consider your child's risk of exposure if they attend public gatherings. In communities with a lot of flu, **people who are at risk of complications from flu should consider staying away from public gatherings.**

If flu is severe in your community, talk to your doctor and child's school to **develop a plan on how to handle your child's special needs.**

Get your child vaccinated for seasonal flu and 2009 H1N1 flu when vaccines are available.

Recognize if your children are sick

Some children may not be able to tell you about their symptoms, which can result in a delay in responding to their illness. It is important to watch carefully for the signs and symptoms of flu or unusual behavior that may be a sign your child is sick. Symptoms of flu include fever, cough, sore throat, runny or stuffy nose, body aches, and fatigue. A fever is a temperature taken with a thermometer that is equal to or greater than 100 degrees Fahrenheit (37.8 degrees Celsius). If you are not able to measure a temperature, your child might have a fever if he or she feels warm, has a flushed appearance, or is sweating or shivering.

Watch for emergency warning signs that need urgent medical attention. These warning signs include:

- Fast breathing or trouble breathing
- Bluish or gray skin color
- Not drinking enough fluids
- Not urinating or no tears when crying
- Severe or persistent vomiting

Not waking up or not interacting
 Being so irritable that the child does not want to be held
 Pain or pressure in the chest or abdomen
 Sudden dizziness
 Confusion
 Flu-like symptoms improve but then return with fever and worse cough

Please share this information with your child’s teacher or other caregivers, so they can tell you if they notice your child is not feeling well.

Tips for taking care of high risk children with the flu

Contact your doctor immediately if your child is sick. This is important because the antiviral medicines used to treat flu work best when started within the first 2 days of getting sick. Your doctor will tell you what special care is needed for your child.

Keep your sick child at home until at least 24 hours after there is no longer a fever or signs of a fever (without the use of a fever-reducing medicine). Keep your child home unless they need to go to the doctor.

Make sure your child gets plenty of rest and drinks clear fluids (such as water, broth, sports drinks, electrolyte beverages for infants, Pedialyte®) to keep from being dehydrated.

If your child has a fever, use fever-reducing medicines that your doctor recommends based on your child’s age. Aspirin (acetylsalicylic acid) **should not** be given to children or teenagers who have flu; this can cause a rare but serious illness called Reye’s syndrome.

Keep your sick child in a separate room (a sick room) in the house as much as possible to limit contact with household members who are not sick. Consider designating a single person as the main caregiver for the sick child.

For more information:

Visit: www.flu.gov
Contact CDC 24 Hours/Every Day
 1-800-CDC-INFO (232-4636)
 TTY: (888)232-6348
cdcinfo@cdc.gov

Page last reviewed August 7, 2009 9:00 AM ET
 Page last updated August 7, 2009 9:00 AM ET
 Content source: [Centers for Disease Control and Prevention](#)

Centers for Disease Control and Prevention 1600 Clifton Rd. Atlanta, GA 30333, USA
 800-CDC-INFO (800-232-4636) TTY: (888) 232-6348, 24 Hours/Every Day - cdcinfo@cdc.gov

